

For sore pain in your shoulders, neck, and low back

# フェイス<sup>®</sup>Z<sub>α</sub> ディクサス<sup>®</sup>ゲル


## Feitas<sup>®</sup>Z<sub>α</sub>Dicsas<sup>®</sup>Gel

Gel product containing 1.0% diclofenac sodium

### Product features

- The active ingredient, diclofenac sodium, is highly effective for relieving pain in the shoulders, low back, joints, and muscle.
- An alcohol gel-based product which is easy-to-apply and dries quick while having a cooling sensation.
- A lightly scented product that you can use without worrying about the people around.
- A milky white gel that is gentle to the skin.

### [Precautions]

 **Do nots** [Failure to follow the instructions may result in worsening symptoms or cause side effects.]

#### 1. Do not use the product if you:

- (1) Have ever experienced allergic symptoms to this product or any ingredient of this product.
- (2) Have ever suffered from asthma.
- (3) Are pregnant or possibly pregnant.
- (4) Children under 15 years old.

#### 2. Do not use the product onto the following areas:

- (1) Areas around the eyes and mucous membranes
- (2) Sensitive areas of the skin (e.g., face, head, armpits)
- (3) Eczema, rash, or wounded areas
- (4) Athlete's foot/ringworm or purulent skin

#### 3. Do not use other external, analgesic, anti-inflammatory drugs while using this product.

#### 4. Avoid long-term, continuous use.

### Consult with a doctor

#### 1. Consult with a doctor, pharmacist, or registered drug seller if you:

- (1) Are receiving treatment at a clinic.
- (2) Have ever experienced allergic symptoms after taking a drug.
- (3) Are receiving the following drug: New quinolone antibiotics

#### 2. Stop use and consult with a doctor, pharmacist, or registered drug seller showing this box if you experience any of the following symptoms after using the product because it may be a side effect:

Affected area	Symptoms
Skin	Eruption/redness, itching, rash, swelling, pain, skin roughness, irritation, pigmentation, blisters, desquamation (fine shedding of skin particles)

In rare cases, the following serious symptoms may occur. In such cases, visit a doctor immediately.

Name of the symptom	Symptoms
Shock (Anaphylaxis)	Skin itching, urticaria, hoarse voice, sneezing, throat itching, suffocating feeling, palpitation, and clouding of consciousness will occur immediately after use.
Contact dermatitis Photosensitivity	Intense symptoms of dermatitis such as eruption/redness accompanied by severe itching, swelling, irritation, and blister/sore, pigmentation, and white spots will appear on the site of application. Eruption/redness, itching, and other symptoms may spread to the whole body in some cases. Symptoms may also develop or worsen in an area exposed to sun light.

#### 3. If the symptoms persist after five or six days of use, stop using the product, and consult with a doctor, pharmacist, or registered drug seller showing this leaflet.

**[Indications]** Low back pain, muscle pain, shoulder pain associated with stiff neck, joint pain, tenosynovitis (hand/wrist pain), elbow pain (e.g., tennis elbow), bruise, sprain

**[Dosage and Usage]** Rub an appropriate amount onto the affected area three to four times a day. Do not cover the applied site with a plastic film or other non-breathing materials. Do not use other external drugs containing the same ingredient as in this product.

<Precautions Related to Dosage and Usage>

- (1) Children under 15 years old should not use this product.
- (2) Strictly follow the dosage and usage.
- (3) Because it is for relieving symptoms such as pain or swelling, but not for treating the disease that causes pain or swelling, use only when you have symptoms.
- (4) Be careful not to apply to the eyes. If it touches the eyes, immediately rinse the eye with cold or lukewarm water. Visit an ophthalmologist if you experience severe symptoms.
- (5) This medicine is only for external use.
- (6) Do not use more than 50 g per week.
- (7) Do not apply other external drugs onto the same area as for this product.
- (8) Do not cover the applied site with a non-breathing materials (e.g., plastic wrap, corrective belt) making it occlusive.
- (9) Wash your hands after use.

**[Ingredients/Contents]**

Ingredients	Content (per 100g)
Diclofenac sodium	1.0g
l-menthol	4.5g
Glycyrrhetic acid	0.05g
Nonylic acid vanillyl amide	0.02g

Contains ethanol, diethanolamine, hydrophobized hydroxypropyl methylcellulose, capsicum tincture, BHT, hydroxypropylcellulose, propylene glycol, polyoxyethylene behenyl ether, and other three ingredients as additives.

**Precautions for Storage and Handling**

- (1) Store in a cool place away from direct sunlight.
- (2) Keep out of children's reach.
- (3) Do not store the product in other containers. (Doing so may lead to misuse or may change its quality.)
- (4) Keep it away from fire.
- (5) Do not let the Gel touch metals such as eye glasses, watches, accessories, clothes made of synthetic material, plastics, and coated surfaces of floors or furniture. The Gel may deteriorate or discolor these items.
- (6) Do not use after their expiration dates.

**Keep it away from flames**

Class 2 petroleum, hazardous grade III, containing ethanol, water soluble

<<Packaging>> 50 g