

Deep stimulation acts on the pain

**サロンパス® EX 温感 インドメタシン 3.5%**

**Salonpas® EX Onkan Indomethacin 3.5%**

<Subtly fragrant> Analgesic antiinflammatory plaster

#### Product features/characteristics

- Contains 3.5% indomethacin and shows excellent analgesic antiinflammatory effect.
- A warming sensation promotes blood circulation in the affected area and alleviates pains induced by stiff neck.
- Adequate sized plaster that is easy for application to areas associated with stiff neck (shoulders, base of the neck, and around the shoulder blades).
- Subtly fragrant plaster that makes you forget that you are wearing it.

#### ⚠[Precautions]

**❌ Do not use:** [Failure to follow directions may result in worsening of current symptoms or cause side effects.]

- Do not use:
  - If you have a history of allergic symptoms to this product or ingredients of this product.
  - If you have a history of asthma.
  - Children under 11 years old.
- Do not use this product on the following areas:
  - Around the eyes and mucous membranes.
  - Eczema, eruptions, and wounds.
  - Athlete's foot/ringworm or purulent skin.

- Do not use this product for more than 2 weeks in a row.



#### Ask a doctor:

- Ask a doctor, pharmacist or registered distributor before use,
  - If you are being treated by a doctor.
  - If you are pregnant or possibly pregnant.
  - If you have a history of allergic symptoms due to drugs.
- If the following symptoms appear after use, they may be side effects. Stop using this product immediately, and ask a doctor, pharmacist, or registered distributor with this box.

Related area	Symptoms
Skin	Rash/redness, itching, swelling, pain, smarting, warmth, dryness, eruption

- If the symptoms do not improve after 5 to 6 days of use, stop using the product and ask a doctor, pharmacist, or registered distributor with this box.

**[Indications]** Stiff shoulder, low back pain, joint pain, muscle pain, tenosynovitis (pain in the hands/wrists), pain in the elbows (tennis elbow), bruises, sprains

**[Dosage and usage]** Remove the plastic film on the surface and apply to the affected area not more than 2 times a day.

<Precautions related to dosage and usage>

- Children under 11 years old should not use this product.
- Children aged 11 and over should use this product only under adult supervision.
- Do not warm the affected area to which this product is applied with a heating pad or electric blanket.
- If you have sensitive skin, apply a small piece (1 to 2 cm square) to the sensitive skin of the inner arm for approximately half a day or longer before application. Use this product only after confirming that the symptoms such as rash/redness, itching, and eruption do not occur.
- Discontinue use at least 1 hour before a bath or shower because it may induce strong stimulation.  
Also, do not use immediately (within 30 minutes) after a bath or shower.
- Clean the skin of the affected area before application.
- Do not apply to the same area continuously if you have sensitive skin.
- If this product is used for bruises or sprains, make sure that swelling has gone down before application.

**[Ingredients/contents]**

Active ingredients	Content (per 100 g mass plaster):
Indomethacin	3.5g
l-menthol	3.5g
Capsicum extract	0.3g

As additional ingredients, this product contains sodium edetate, cycloaliphatic saturated hydrocarbon resin, styrene-isoprene-styrene block copolymer, BHT, polyisobutylene, and liquid paraffin, as well as 2 other ingredients.

**Precautions for storage and handling**

- (1) Store in a cool place away from direct sunlight.
- (2) Keep out of reach of children.
- (3) Avoid storing the product in other containers (it may result in a misuse or a change in quality).
- (4) Place unused products in the pouch and fold the opening properly for storage.
- (5) Do not use the product after the expiration date.

<Packaging> 5.4cm × 8.56cm .... 20 patches, 40 patches