

Directly acts on sore pain

インドメタシン(0.5%配合)

# サロシップ® インドメタシン EX

## Salonsip® Indomethacin EX

インドメタシン(0.5%)

&lt;Cooling&gt; Analgesic antiinflammatory gel patch

**Product features/characteristics**

- Contains indomethacin and shows excellent analgesic antiinflammatory effect.  
“Indomethacin” that directly acts on the cause of persistent pain in the lower back and joints gradually penetrates into the skin and shows excellent analgesic antiinflammatory effects.
- Moisture-keeping zipper  
Inner pouch has a moisture-keeping zipper to protect the gel patches from drying, making it convenient for storage.

**⚠[Precautions]****❌ Do not use:** [Failure to follow directions may result in worsening of current symptoms or cause side effects.]

- Do not use:
  - If you have a history of allergic symptoms to this product or ingredients of this product.
  - If you have a history of asthma.
  - Children under 15 years old.
- Do not use this product on the following areas:
  - Around the eyes and mucous membranes.
  - Eczema, eruptions, and wounds.
  - Athlete's foot/ringworm or purulent skin.
- Do not use this product for more than 2 weeks in a row.

**Ask a doctor:**

- Ask a doctor, pharmacist or registered distributor before use,
  - If you are being treated by a doctor.
  - If you are pregnant or possibly pregnant.
  - If you have a history of allergic symptoms due to drugs.
- If the following symptoms appear after use, there may be side effects. Stop using this product immediately, and ask a doctor, pharmacist, or registered distributor with this box.

Related area	Symptoms
Skin	Rash/redness, itching, swelling, smarting, warmth, dryness

- If the symptoms do not improve after 5 to 6 days of use, stop using the product and ask a doctor, pharmacist, or registered distributor within this box.

**[Indications]** Joint pain, tenosynovitis (pain in the hands/wrists), pain in the elbows (tennis elbow), muscle pain, low back pain, bruises, sprains, shoulder pain associated with stiff neck

**[Dosage and usage]** Remove the plastic film and apply to the affected area, not more than twice a day.

<Precautions related to dosage and usage>

- Children under 15 years old should not use this product.
- This product only treats symptoms such as pain and swelling that are caused by a disease/illness, but not the disease/illness itself. Use this product only when symptoms occur.
- Wipe and clean the affected area thoroughly when sweating or if it is wet.
- If you have sensitive skin, apply a small piece (1 to 2 cm square) to the sensitive skin of the inner arm for approximately half a day or longer before full application. Use this product only after confirming that the symptoms such as rash/redness, itching, and eruption do not occur.

**[Ingredients/contents]** Per 100 g mass plaster: Indomethacin.....0.5 g

As additional ingredients, this product contains sodium edetate, N-methyl-2-pyrrolidone, l-menthol, kaolin, carboxyvinyl polymer, sodium carboxymethylcellulose, glycerin, magnesium aluminosilicate, titanium oxide, D-sorbitol, partially neutralized polyacrylic acid, polyoxyethylene hydrogenated castor oil, polyvinyl alcohol (partial saponification product), and water.

**Precautions for storage and handling**

- (1) Store in a cool dry place away from direct sunlight.
- (2) Keep out of reach of children.
- (3) Avoid storing the product in other containers (it may result in a misuse or a change in quality).
- (4) Place the unused products in the pouch and zip up or fold the opening for storage.
- (5) Do not use the product after the expiration date.

<Packaging> 10cm × 14cm .... 12 patches, 24 patches