

For sore pain



Feitas[®] Z_α Dicsas[®] Oban

Contains 2.0% diclofenac sodium/l-menthol

Product features/characteristics

- Transdermal analgesic antiinflammatory tape with 2.0% diclofenac sodium, an active ingredient.
- Contains 3.5% l-menthol for enhanced feeling of effectiveness.
- Shows excellent effects on sore pain in the shoulders, neck, lower back, joints, and muscles.
- Subtly fragrant product you can use comfortably during bedtime or in public.
- Stretches to all directions for the perfect fit on the skin.
- Large sized patch that can cover wide area such as the lower back with 1 patch.

⚠️ [Precautions]

❌ Do not use: [Failure to follow directions may result in worsening of current symptoms or cause side effects.]

- Do not use:
 - (1) If you have a history of allergic symptoms to this product or ingredients of this product.
 - (2) If you have a history of asthma.
 - (3) If you are pregnant or possibly pregnant.
 - (4) Children under 15 years old.
- Do not use this product on the following areas:
 - (1) Around the eyes and mucous membranes.
 - (2) Eczema, eruptions, and wounds.
 - (3) Athlete's foot/ringworm or purulent skin.
- Do not use this product with, or at the same time as, other external analgesic antiinflammatory products.
- Do not use this product for more than 2 weeks in a row.



Ask a doctor:

- Ask a doctor, pharmacist, or registered distributor before use,
 - (1) If you are being treated by a doctor.
 - (2) If you are using other pharmaceutical product.
 - (3) If you have a history of allergic symptoms due to drugs.
 - (4) If you have a history of eruption induced by a tape.
 - (5) If you have been diagnosed with the following: Peptic ulcer, hematologic disorder, liver disorder, renal disorder, hypertension, cardiac disorder, influenza
 - (6) If you are receiving the following drugs: New quinolone antibiotics, triamterene, lithium, methotrexate, nonsteroidal antiinflammatory analgesics, steroids, diuretics, cyclosporine, selective serotonin reuptake inhibitors
 - (7) If you are an elderly person.
- If the following symptoms appear after use, there may be side effects. Stop using this product immediately, and ask a doctor, pharmacist, or registered sales representative with this box.

Related area	Symptoms
Skin	Rash/redness, itching, eruption, swelling, skin roughness, irritation, pigmentation, blisters, desquamation (fine shedding of skin particles)

In rare cases, the following serious symptoms may occur. In such cases, consult a doctor immediately.

Name of the symptom	Symptoms
Shock (anaphylaxis)	Skin itching, urticaria, hoarse voice, sneezing, throat itching, feeling of dyspnea, palpitation, and clouding of consciousness occur immediately after use.
Contact dermatitis Photosensitivity	Intense symptoms of dermatitis such as rash/redness, swelling, irritation, and blisters/sores that are accompanied by intense itching, pigmentation, and white spots appear on the site of application. In some cases, symptoms such as rash/redness and itching may spread all over the body. Also, symptoms may appear or worsen after the relevant area comes into contact with the sun light.

- If the symptoms persist after 5 to 6 days of use, stop using the product and ask a doctor, pharmacist, or registered distributor with this box.

[Indications] Low back pain, muscle pain, shoulder pain associated with stiff shoulder, joint pain, tenosynovitis (pain in the hands/wrists), pain in the elbows (tennis elbow), bruises, sprains

[Dosage and usage] Remove the plastic film and apply 1 patch to the affected area once daily. Do not use more than 1 patch per application. Also, do not use with, or at the same time as, other external application products that contain the ingredients of this product.

<Precautions related to dosage and usage>

- (1) Children under 15 years old should not use this product.
- (2) Do not use otherwise than as specified in the dosage and usage.
- (3) Since this product only treats symptoms such as pain and swelling, rather than the disease that caused it, use this product only when symptoms occur.
- (4) If you have sensitive skin, apply a small piece (1 to 2 cm square) to the sensitive skin of the inner arm for approximately half a day or longer before application. Use this product only after confirming that the symptoms such as rash/redness, itching, and eruption are not induced.
- (5) Avoid leaving a patch continuously for over 24 hours. If applying a new patch to the same affected area, confirm that symptoms such as rash/redness, itching, and eruption do not occur.
- (6) Do not use with, or at the same time as, other external application products.
- (7) Wipe and clean the affected area thoroughly when sweating or if it is wet.

[Ingredients/contents]

Active ingredients	Content (per 100 g mass plaster):
Diclofenac sodium	2.0g
l-menthol	3.5g

As additional ingredients, this product contains styrene-isoprene-styrene block copolymer, terpene resin, BHT, polyisobutylene, and liquid paraffin, as well as 2 other ingredients.

Precautions for storage and handling

- (1) Store in a cool dry place away from direct sunlight.
- (2) Keep out of reach of children.
- (3) Avoid storing the product in other containers (it may result in a misuse or a change in quality).
- (4) Do not use the product after the expiration date.
- (5) Place unused products in the pouch and fold the opening properly for storage.

<Packaging> 10cm × 14cm 7 patches