

Feitas_®Shippu Onkan

Gel patch with felbinac

Product features/characteristics

- Analgesic antiinflammatory gel patch containing 70 mg per patch of "Felbinac" with excellent antiinflammatory/analgesic effect.
- "Felbinac" inhibits the production of a substance involved in pain (prostaglandin) and shows excellent effects on pain in the lower back, shoulders, joints, and muscles.
- Soothing warming sensation of capsicum extract improves blood circulation and acts well on shoulder pain, low back pain, and joint pain.
- Round corners prevent stucking and peeling by clothing.
- Freely stretchable patch that fits the joint area perfectly.
- "Baripita sheet" with a film that can be removed by simply pulling the patch to the left and right for easy application to any site.

· /t [Precautions]

Do not use: [Failure to follow directions may result in worsening of current symptoms or cause side effects.]

- 1. Do not use:
 - (1) If you have a history of allergic symptoms to this product or ingredients of this product.
 - (2) If you have a history of asthma.
 - (3) If you are pregnant or possibly pregnant.
 - (4) Children under 15 years old.
- 2. Do not use this product on the following areas:
 - (1) Around the eyes and mucous membranes.
 - (2) Eczema, eruptions, and wounds.
 - (3) Athlete's foot/ringworm or purulent skin.
- 3. Do not use this product for more than 2 weeks in a row.



- 1. Ask a doctor, pharmacist, or registered sales representative before use,
 - (1) If you are being treated by a doctor.
 - (2) If you have a history of allergic symptoms due to drugs.
- 2. If the following symptoms appear after use, they may be side effects. Stop using this product immediately, and ask a doctor, pharmacist, or registered distributor with this box.

Related area	Symptoms	
Skin	Rash/redness, swelling, pain, itching, smarting, eruption	

In rare cases, the following serious symptoms may occur. In such cases, consult a doctor immediately.

Name of the symptom	Symptoms	
Shock	Skin itching, urticaria, hoarse voice, sneezing, throat itching, feeling of dyspnea,	
(anaphylaxis)	palpitation, and clouding of consciousness occur immediately after use.	

3. If the symptoms persist after 5 to 6 days of use, stop using the product and ask a doctor, pharmacist, or registered distributor with this box.

[Indications] Shoulder pain associated with stiff shoulder, low back pain, joint pain, muscle pain, tenosynovitis (pain and swelling in the hands, wrists, and ankles), pain in the elbows (tennis elbow), bruises, sprains

[Dosage and usage] Apply to the affected area no more than 2 times a day.

- <Pre><Precautions related to dosage and usage>
 - (1) Children under 15 years old should not use this product.
 - (2) Do not use otherwise than as specified in the dosage and usage.
 - (3) Since this product only treats symptoms such as pain and swelling, rather than the disease itself, use this product only when there are symptoms.
 - (4) Do not warm the affected area to which this product is applied with a heating pad or electric blanket.
 - (5) Discontinue use at least 1 hour before a bath or shower because it may cause strong stimulation. Also, do not use immediately (within 30 minutes) after a bath or shower.
 - (6) Clean the skin of the affected area before application.
 - (7) Do not apply to the same area continuously if you have sensitive skin.
 - (8) If this product is used for bruises or sprains, make sure that swelling has gone down before application.

[Ingredients/contents]

Active ingredients	Content (per 100 g mass plaster):
Felbinac	0.7g
Capsicum extract	0.1 g (1.25 g as raw herbal medicine)

As additional ingredients, this product contains sodium edetate, N-methyl-2-pyrrolidone, I-menthol, glycerin, silicate AI, titanium oxide, tartrate, gelatin, partially neutralized polyacrylic acid, polyvinyl alcohol (partial saponification product), and magnesium aluminometasilicate, as well as 3 other ingredients.

Precautions for storage and handling

- (1) Store in a cool dry place away from direct sunlight.
- (2) Keep out of reach of children.
- (3) Avoid storing the product in other containers (it may result in a misuse or a change in quality).
- (4) Do not use the product after the expiration date.
- (5) Once the package is opened, zip up the pouch tightly for storage.

<Packaging>10cm × 14cm 16 patches