

For sore pain in the muscles/joints

エアーサロンパスDX

フェルビナク
3.0%配合

Air[®]Salonpas[®]DX


Contains 3.0% felbinac

Analgesic antiinflammatory spray

Product features/characteristics

- Analgesic antiinflammatory spray with 3.0% “Felbinac,” an ingredient that has been demonstrated to show excellent antiinflammatory analgesic effect and 3.0% “l-menthol,” an ingredient that improves blood circulation.
- When sprayed onto the affected area such as joint pain, muscle pain, bruises, and sprains, the active ingredient “Felbinac” is absorbed through the skin and alleviates sore pain, in addition to the cooling effect.

[Precautions]

 **Do not use:** [Failure to follow directions may result in worsening of current symptoms or cause side effects.]

1. Do not use:

- (1) If you have a history of allergic symptoms to this product or ingredients of this product.
- (2) If you have a history of asthma.
- (3) If you are pregnant or possibly pregnant.
- (4) Children under 15 years old.

2. Do not use this product on the following areas:

- (1) Around the eyes and mucous membranes.
- (2) Eczema, eruptions, and wounds.
- (3) Purulent areas.



Ask a doctor:

1. Ask a doctor, pharmacist, or registered distributor before use,

- (1) If you are being treated by a doctor.
- (2) If you have a history of allergic symptoms due to drugs.

2. If the following symptoms appear after use, they may be side effects. Stop using this product immediately, and ask a doctor, pharmacist, or registered distributor with this information leaflet.

Related area	Symptoms
Skin	Rash/redness, swelling, itching, smarting, eruption

In rare cases, the following serious symptoms may occur. In such cases, consult a doctor immediately.

Name of the symptom	Symptoms
Shock (anaphylaxis)	Skin itching, urticaria, hoarse voice, sneezing, throat itching, feeling of dyspnea, palpitation, and clouding of consciousness occur immediately after use.


3. If the symptoms persist after about a week of use, stop using the product and ask a doctor, pharmacist, or registered distributor with this information leaflet.

[Indications] Stiff shoulder, low back pain, joint pain, muscle pain, bruises, sprains, tenosynovitis (pain and swellings in the hands, wrists, and ankles), pain in the elbows (tennis elbow)

[Dosage and usage] Spray or apply a proper amount on the affected area 2 to 4 times a day.

<Precautions related to dosage and usage>

- (1) Children under 15 years old should not use this product.
- (2) Do not use otherwise than as specified in the dosage and usage.
- (3) Avoid contact with the eyes. If eye contact occurs, rinse immediately with water or lukewarm water.
Consult an ophthalmologist if the symptoms are severe.
- (4) Do not spray toward the face.
- (5) Do not use this product in large amount or on large area.
- (6) For external use only. Avoid inhalation. Inhalation may result in symptoms of irritation to the nose and throat such as dizziness, nausea, and cough. Avoid inhalation as much as possible and pay attention to the surrounding people when using this product.
- (7) Shake the can well and always spray with the nozzle pointing downward (if the nozzle is pointing upward or sideways, only the gas will come out; the drug solution will remain in the can and you will not be able to use it up completely or the drug solution may splash).
- (8) Hold the can 10 cm away from the affected area to spray.
- (9) Spray each affected area for no longer than 3 seconds (it may induce frostbite).
- (10) If you are going to apply this on the neck or shoulder, spray it to your hand first and rub your neck/shoulder with your hand.
- (11) Do not cover the affected area with unbreathable materials such as plastic wrap film.



[Ingredients/contents]

Active ingredients	Content (per 100 g stock solution):
Felbinac	3g
l-menthol	3g

As additional ingredients, this product contains ethanol, triethanolamine, hydroxypropylcellulose, polyoxyethylene hydrogenated castor oil, and DME.

Precautions for storage and handling

- (1) Store in a cool dry place with cap closed tightly, away from direct sunlight.
- (2) Keep out of reach of children.
- (3) Avoid storing the product in other containers (it may result in a misuse or a change in quality).
- (4) Be careful not to let this product come into contact with clothing, furnitures, leather products, or chemical products.
It may deteriorate or discolor.
- (5) Keep away from fire.
- (6) Do not use this product in places with fire sparks.

<Packaging> 80mL, 120mL

[Precautions and method of degassing for disposal]

<Precautions for degassing>

- Make sure to perform the following procedures outdoor away from fire after using up the contents completely.
- During degassing, turn your face away from the nozzle because the slightly remaining drug solution comes out together with the gas.
- Do not use those other than the cap of this product for degassing.
- Be careful not to directly inhale the gas as it comes out from the can.

<How to degas>

* The figure below is an information leaflet of the product. Sample product has a different structure.

- (1) Press "PUSH" mark to remove the button and shoulder cover.
- (2) Pull out the push button.
- (3) Remove the upper part of the cap and place it up-side down on the flat surface.
- (4) Push the main body into the hole in the center of the top face of the cap and hold it until the hissing sound is no longer heard.
- (5) Dispose according to local guidelines after the hissing sound is no longer heard.

