

Fast-drying, no-mess, liquid formula for pain in the muscles/joints

フェイスローション 微香性

Feitas[®] Lotion Slightly fragrant

Percutaneous analgesic antiinflammatory agent

Product features/characteristics

- Feitas Lotion is a transdermal analgesic antiinflammatory agent with “Felbinac” that has been demonstrated to show excellent antiinflammatory/analgesic effect as the primary ingredient (3%).
- “Felbinac” inhibits the production of a substance involved in pain (prostaglandin) and shows excellent effects on pain in the shoulders, lower back, joints, and muscles.
- Contains 3.0% l-menthol for refreshing feeling of use.
- Ultimate easy-to-hold and easy-to-apply original bottle for no-mess, easy application to the shoulders and neck by yourself.
- Subtly fragrant product you can use without worrying about the surrounding people.

[Precautions]

 **Do not use:** [Failure to follow directions may result in worsening of current symptoms or cause side effects.]

1. Do not use:

- (1) If you have a history of allergic symptoms to this product or ingredients of this product.
- (2) If you have a history of asthma.
- (3) If you are pregnant or possibly pregnant.
- (4) Children under 15 years old.

2. Do not use this product on the following areas:

- (1) Around the eyes and mucous membranes.
- (2) Eczema, eruptions, and wounds.
- (3) Purulent areas.



Ask a doctor:

1. Ask a doctor, pharmacist, or registered distributor before use,

- (1) If you are being treated by a doctor.
- (2) If you have a history of allergic symptoms due to drugs.

2. If the following symptoms appear after use, they may be adverse reactions. Stop using this product immediately, and ask a doctor, pharmacist, or registered distributor with this information leaflet.

Related area	Symptoms
Skin	Rash/redness, swelling, itching, smarting, eruption

In rare cases, the following serious symptoms may occur. In such cases, consult a doctor immediately.

Name of the symptom	Symptoms
Shock (anaphylaxis)	Skin itching, urticaria, hoarse voice, sneezing, throat itching, feeling of dyspnea, palpitation, and clouding of consciousness occur immediately after use.

3. If the symptoms persist after about a week of use, stop using the product and ask a doctor, pharmacist, or registered distributor with this information leaflet.

[Indications] Shoulder pain associated with stiff shoulder, low back pain, joint pain, muscle pain, bruises, sprains, tenosynovitis (pain and swellings in the hands, wrists, and ankles), pain in the elbows (tennis elbow)

[Dosage and usage] Apply a proper amount to the affected area 2 to 4 times a day.

<Precautions related to dosage and usage>

- (1) Children under 15 years old should not use this product.
- (2) Do not use otherwise than as specified in the dosage and usage.
- (3) Avoid contact with the eyes. If eye contact occurs, rinse immediately with water or lukewarm water.
Consult an ophthalmologist if the symptoms are severe.
- (4) For external use only.
- (5) Do not cover the affected area with unbreathable materials such as plastic wrap film after application of the drug.

[Ingredients/contents]

Active ingredients	Content (per 100 g):	Functions
Felbinac	3.0g	Suppresses inflammation and alleviates pain.
l-menthol	3.0g	Provides a cooling sensation and alleviates pain.

As additional ingredients, this product contains ethanol, N-methyl-2-pyrrolidone, triethanolamine, octaacetyl sucrose, benzyl alcohol, polyoxyethylene hydrogenated castor oil, polyoxyethylene polyoxypropylene cetyl ether, and isopropyl myristate.

Precautions for storage and handling

- (1) Store in a cool dry place with cap closed tightly, away from direct sunlight.
- (2) Keep out of reach of children.
- (3) Avoid storing the product in other containers (it may result in a misuse or a change in quality).
- (4) Keep away from fire.
- (5) Metals such as glasses, watches, and accessories, clothing made of synthetic fibers, plastics, and painted surfaces of the floors and furnitures may deteriorate or be discolored when they come into contact with this product. Make sure that this product does not come into contact with these materials.
- (6) Do not use the product after the expiration date. Also, use as soon as possible after opening the package even if the expiration date has not passed.

<Packaging> 50mL