

For sore pain in the lower back/knees




Feitas[®] Z Cream Slightly
fragrant

Contains 1.0% diclofenac sodium

Product features/characteristics

- An active ingredient, diclofenac sodium shows excellent effects on sore pain in the shoulders, lower back, and joints.
- Smooth, less sticky cream product that spreads well even in small amount.
- Subtly fragrant product you can use during bedtime or in public.
- Contained in a laminated tube such that you can use it up completely.

[Precautions]

 **Do not use:** [Failure to follow directions may result in worsening of current symptoms or cause side effects.]

- Do not use:
 - If you have a history of allergic symptoms to this product or ingredients of this product.
 - If you have a history of asthma.
 - If you are pregnant or possibly pregnant.
 - Children under 15 years old.
- Do not use this product on the following areas:
 - Around the eyes and mucous membranes.
 - Sensitive areas of the skin (face, head, armpits).
 - Eczema, eruptions, and wounds.
 - Athlete's foot/ringworm or purulent skin.
- Do not use this product with, or at the same time as, other external analgesic antiinflammatory products.
- Avoid long-term, continuous use.



Ask a doctor:

- Ask a doctor, pharmacist or registered distributor before use,
 - If you are being treated by a doctor.
 - If you have a history of allergic symptoms due to drugs.
 - If you are receiving the following drugs: New quinolone antibiotics
- If the following symptoms appear during or after use, they may be side effects. Stop using this product immediately, and ask a doctor, pharmacist, or registered distributor with this information leaflet.

Related area	Symptoms
Skin	Rash/redness, itching, eruption, swelling, pain, irritation, warmth, skin roughness, desquamation (peeled skin like scurf or dirt), blisters, pigmentation

In rare cases, the following serious symptoms may occur. In such cases, consult a doctor immediately.

Name of the symptom	Symptoms
Shock (anaphylaxis)	Skin itching, urticaria, hoarse voice, sneezing, throat itching, feeling of dyspnea, palpitation, and clouding of consciousness occur immediately after use.
Contact dermatitis Photosensitivity	Intense symptoms of dermatitis such as rash/redness, swelling, irritation, and blisters/sores that are accompanied by intense itching, pigmentation, and white spots appear on the site of application. In some cases, symptoms such as rash/redness and itching may spread all over the body. Also, symptoms may appear or worsen after the relevant area comes into contact with the sun light.

- If the symptoms persist after 5 to 6 days of use, stop using the product and ask a doctor, pharmacist, or registered distributor with this information leaflet.

[Indications] Low back pain, shoulder pain associated with stiff shoulder, joint pain, muscle pain, tenosynovitis (pain in the hands/wrists), pain in the elbows (tennis elbow), bruises, sprains

[Dosage and usage] Rub a proper amount to the affected area 3 to 4 times a day. However, do not cover the site of application with unbreathable materials such as plastic wrap film. Also, do not use with, or at the same time as, other external application products that contain the ingredients of this product.

<Precautions related to dosage and usage>

- (1) Do not use otherwise than as specified in the dosage and usage.
- (2) For external use only. Do not take it orally.
- (3) Do not use more than 50 g per week.
- (4) Avoid contact with the eyes. If eye contact occurs, rinse immediately with water or lukewarm water. Consult an ophthalmologist if the symptoms are severe.
- (5) Do not cover the affected area with unbreathable materials such as plastic wrap film after rubbing this product.
- (6) Wash hands after use.

[Ingredients/contents]

Active ingredients	Content (per 1 g):
Diclofenac sodium	10mg

As inactive ingredients, this product contains l-menthol, white petrolatum, isopropyl myristate, squalane, propylene glycol, polyoxyethylene hydrogenated castor oil, dibutyl hydroxy toluene, methyl-p-hydroxybenzoate, butyl-p-hydroxybenzoate, carboxyvinyl polymer, and pH regulator, as well as 2 other ingredients.

Precautions for storage and handling

- (1) Store in a cool place with lid closed tightly, away from direct sunlight.
- (2) Keep out of reach of children.
- (3) Avoid storing the product in other containers (it may result in a misuse or a change in quality).
- (4) Do not use the product after the expiration date (specified on the paper box and container). Also, use as soon as possible after opening the package even if the expiration date has not passed for maintenance of quality.

<Packaging> 30g