

For sore pain in the neck/shoulders




Feitas[®] Z_α Lotion Slightly fragrant

Contains 1.0% diclofenac sodium

Product features/characteristics

- An active ingredient, diclofenac sodium shows excellent effects on sore pain in the shoulders, lower back, and joints.
- Contains 3.0% l-menthol for refreshing feeling of use.
- The ultimate easy-to-hold and easy-to-apply original bottle for no-mess, easy application to the shoulders and neck by yourself.
- Smell-free, subtly fragrant product you can use without worrying about the surrounding people.

[Precautions]

 **Do not use:** [Failure to follow directions may result in worsening of current symptoms or cause side effects.]

1. Do not use:

- (1) If you have a history of allergic symptoms to this product or ingredients of this product.
- (2) If you have a history of asthma.
- (3) If you are pregnant or possibly pregnant.
- (4) Children under 15 years old.

2. Do not use this product on the following areas:

- (1) Around the eyes and mucous membranes.
- (2) Sensitive areas of the skin (face, head, armpits).
- (3) Eczema, eruptions, and wounds.
- (4) Athlete's foot/ringworm or purulent skin.

3. Do not use this product with, or at the same time as, other external analgesic antiinflammatory products.

4. Avoid long-term, continuous use.



Ask a doctor:

1. Ask a doctor, pharmacist, or registered sales representative distributor before use,

- (1) If you are being treated by a doctor.
- (2) If you have a history of allergic symptoms due to drugs.
- (3) If you are receiving the following drugs: New quinolone antibiotics

2. If the following symptoms appear after use, there may be side effects. Stop using this product immediately, and ask a doctor, pharmacist, or registered sales representative distributor with this information leaflet.

Related area	Symptoms
Skin	Rash/redness, itching, eruption, swelling, skin roughness, irritation, pigmentation, blisters, desquamation (fine shedding of skin particles)

In rare cases, the following serious symptoms may occur. In such cases, consult a doctor immediately.

Name of the symptom	Symptoms
Shock (anaphylaxis)	Skin itching, urticaria, hoarse voice, sneezing, throat itching, feeling of dyspnea, palpitation, and clouding of consciousness occur immediately after use.
Contact dermatitis Photosensitivity	Intense symptoms of dermatitis such as rash/redness, swelling, irritation, and blisters/sores that are accompanied by intense itching, pigmentation, and white spots appear on the site of application. In some cases, symptoms such as rash/redness and itching may spread all over the body. Also, symptoms may appear or worsen after the relevant area comes into contact with the sun light.

3. If the symptoms persist after 5 to 6 days of use, stop using the product and ask a doctor, pharmacist, or registered sales representative distributor with this information leaflet.

[Indications] Low back pain, muscle pain, shoulder pain associated with stiff shoulder, joint pain, tenosynovitis (pain in the hands/wrists), pain in the elbows (tennis elbow), bruises, sprains

[Dosage and usage] Apply a proper amount to the affected area 3 to 4 times a day. However, do not cover the site of application with unbreathable materials such as plastic wrap film. Also, do not use with, or at the same time as, other external application products that contain the ingredients of this product.

<Precautions regarding to dosage and usage>

- (1) Children under 15 years old should not use this product.
- (2) Do not use other than as specified in the dosage and usage.
- (3) Since this product only treats symptoms such as pain and swelling, rather than the disease that caused it, use this product only when symptoms occur.
- (4) For external use only.
- (5) Do not use more than 50 mL per week.
- (6) Avoid contact with the eyes. If eye contact occurs, rinse immediately with water or lukewarm water. Consult an ophthalmologist if the symptoms are severe.
- (7) Do not use with, or at the same time as, other external application products.
- (8) Do not cover the site of application with unbreathable materials (e.g., plastic wrap film and correction belt) that induces a sealed condition.

[Ingredients/contents]

Active ingredients	Content (per 100 g):
Diclofenac sodium	1g
l-menthol	3g

As additional ingredients, this product contains diisopropyl adipate, isopropanol, lactate, propylene glycol, and polysorbate 80.

Precautions for storage and handling

- (1) Store in a cool place away from direct sunlight.
- (2) Keep out of reach of children.
- (3) Avoid storing the product in other containers (it may result in a misuse or a change in quality).
- (4) Keep away from fire.
- (5) Metals such as glasses, watches, and accessories, clothing made of synthetic fibers, plastics, and painted surfaces of the floors and furnitures may deteriorate or be discolored when they come into contact with this product. Make sure that this product does not come into contact with these materials.
- (6) Do not use the product after the expiration date.

Keep away from fire

Class II petroleum, water-soluble, hazard class III (isopropanol, propylene glycol)

<Packaging> 50mL